Post-operative Instructions for Wart Procedures

Doug Ichikawa, D.P.M.

Suzanne Wilson, D.P.M.

Now that your surgery has been completed, you can greatly enhance the outcome by following these instructions.

- 1. Please call us! Call us if you have any questions or concerns regarding your surgery or aftercare of the surgery site.
- 2. Go directly home and keep your foot as elevated as possible on the way home.
- 3. At home, **keep your foot elevated** as much as possible. it should be a few inches above the level of your heart.
- 4. **Keep your dressings dry and clean for 3 days**. After 3 days begin soaking your foot as described below.
- 5. A small amount of bleeding into the bandage is very normal and should cause no alarm. If you have active or excessive bleeding, please call your doctor.
- 6. Mild to moderate discomfort in the area of the surgery can be expected. This is best treated by elevation, ice and taking **tylenol**. If your pain is not controlled by these measures, please call your doctor.
- 7. Please **refrain** from alcoholic beverages, smoking, Bungee jumping and the operation of bulldozers or other heavy machinery!
- 8. Your doctor would like to see you for your first **postop visit within the first week** after surgery. Please call to make this appointment.

Again, please call us if you have any questions or concerns about your surgery. Our doctors are on call 24 hours a day to assist you in cases of emergency.

Soaking Instructions:

- 1. Fill a shallow pan with warm water.
- 2. Soak your foot for about **5-10 minutes** .
- (do this with the dressing on the first time so that it will come off much easier.)
- 3. Gently dry your foot and apply an antibiotic cream or ointment.
- 4. Apply a clean bandaid or gauze dressing
- 5. Repeat the soaking **once a day** until for at least two weeks.
- 6. Hydrogen peroxide may be use to gently scrub and cleanse the excision area.

Bellevue Podiatric Physicians

(425) 283-5093